arXiv:1211.0501v1 [physics.data-an] 1 Nov 2012

Surprisingly Rational: Evidence that people follow probability theory when judging probabilities, and that biases in judgment are due to noise.

Fintan Costello,
School of Computer Science and Informatics,
University College Dublin,
Belfield, Dublin 4, Ireland,

Paul Watts
Department of Mathematical Physics,
National University of Ireland Maynooth,
Maynooth, Co Kildare, Ireland

Abstract

The systematic biases and errors seen in people's probability judgments are typically taken as evidence that people do not reason about probability using the rules of probability theory. We show the contrary: that these biases are a consequence of people correctly following probability theory, but with random variation or noise affecting the reasoning process. Taking $P_E(A)$ to represent a person's estimate for the probability of some event A, this random variation account predicts that on average $P_E(A) + P_E(B) - P_E(A \vee B) - P_E(A \wedge B) = 0$ for all pairs of events A, B, just as required by probability theory. Analysing data from an experiment asking people to estimate such probabilities for a number of pairs A, B we find striking confirmation of this prediction.

1 Introduction

The ability to reason with uncertain knowledge (that is, to reason with probabilities) is crucial to the ability to survive and prosper in "an ecology that is of essence only partly accessible to foresight" (Brunswik, 1955). It is therefore reasonable to expect that humans, having prospered in such an ecology, would be able to reason about probabilities extremely well: any ancestors who could not reason effectively about probabilities would not survive long, and so the biological basis of their reasoning would be driven from the gene pool. Probability theory provides a calculus of chance describing how to make optimal predictions under uncertainty: taking the argument one step further, it is reasonable to expect that our probabilistic reasoning will follow the rules of probability theory.

The conventional view in current psychology is that this expectation is wrong. Instead, the dominant position is that

> In making predictions and judgments under uncertainty, people do not appear to follow the calculus of chance or the statistical theory of prediction. Instead they rely on a limited number of heuristics which sometimes yield reasonable judgments and sometimes lead to severe and systematic errors (Tversky and Kahneman, 1973, p. 237)

This conclusion is based on series of systematic and reliable biases in people's judgements of probability, many identified in the 1970's and 1980's by Tversky, Kahneman and colleagues. The heuristics and biases approach has reached a level of popularity rarely seen in psychology, with Kahneman recieving a Nobel Prize for his work in this area, and with numerous popular psychology books and newspaper articles presenting this view to the general public. Indeed, the idea that people cannot reason with probabilities has become a widespread truism: for example, the Science Gallery in Dublin recently announced an exhibition on risk which it describes as "enabling visitors to explore our inability to determine the probability of everything from a car crash to a coin toss" (The Irish Times - Thursday, October 11, 2012).

In this paper we describe a very simple alternative to the heuristics and biases view. In this account people reason about probability according to the rules of probability theory, but are subject to random variation or noise in this reasoning process. While it may seem that noise in the workings of a rational system will result in "nothing more than error variance centered around a normative response"(Shafir and Leboeuf, 2002), in fact there are a number of ways in which random variation can produce systematic and reliable deviations from the normatively correct responses. These systematic deviations often correspond to the observed patterns of bias seen in people's probability judgments.

This 'probability theory plus noise' account of biases in human probabilistic reasoning has a critical characteristic that differentiates it from the heuristicsaccount: the random variation account predicts that there should, on average, be no systematic bias or error in people's judgements for some specific probabilities. Taking $P(A)$ to represent the objective, true probability of some event A, $P_E(A)$ to represent a reasoner's estimate of that probability, and $\overline{P_E}(A)$ to represent the mean or expected value of $P_E(A)$ (the average estimate of the probability of event A), this random variation account predicts that

$$\overline{P_E}(A)+\overline{P_E}(B)-\overline{P_E}(A\wedge B)-\overline{P_E}(A\vee B) = P(A)+P(B)-P(A\wedge B)-P(A\vee B) = 0$$

for all A and B. This prediction follows directly from the familiar 'addition law' of probability theory, which states that

$$P(A \vee B) = P(A) + P(B) - P(A \wedge B) \tag{1}$$

(because the size of the set of events $A \vee B$ is equal to the size of the set of events A plus the size of the set B, minus the union of those two sets). Analysing data from an experiment asking people to estimate such probabilities for a number of pairs A, B we find strikingly strong confirmation of this prediction, showing that people's probability judgments embody the addition law.

The organisation of the paper is as follows. In the first section we present our very simple account and show how it can explain well-established and systematic errors in human probabilistic reasoning: conservatism, subadditivity, the conjunction error, and the disjunction error. In the second section we describe our experimental results confirming the above prediction, and in the final section discuss some implications of this result.

2 The systematic influence of noise

In this section we present our simple account of the influence of noise on people's probability estimates by showing how this account can explain 4 well-known patterns of bias in people's probability judgment: conservatism or underconfidence, subadditivity, the conjunction error, and the disjunction error. Other more complex accounts also based on noise have previously been given for each of these biases (see Costello, 2009a,b, Erev et al., 1994, Hilbert, 2012): as far as we know this is the first account explaining all 4 biases, and the first to make specific predictions about when bias will vanish.

In discussing the influence of random variation on probabilistic reasoning we assume a rational reasoner with a long-term episodic memory. For simplicity we assume that, apart from random variation, the reasoner is 'perfect': that is, if the reasoner were not subject to random variation then each estimate $P_E(A)$ would be equal to the correct probability $P(A)$. We assume a simple form of long-term memory containing n episodes where each recorded episode i contains a flag $f_i(A)$ that is set to 1 if i contains event A and set to 0 otherwise, and the reasoner estimates the probability of event A by counting these flags. We assume a minimal form of transient error, in which there is some small probability d that when the state of some flag $f_i(A)$ is read, the value obtained is not the correct value for that flag. We take $C(A)$ to be number of flags that were read as 1 in some particular query of memory, and T_A be the number of flags whose correct value is actually 1. Note that due to random error $C(A) = T_A$ does not necessarily hold.

2.1 Conservatism

Probabilities range in value between 0 and 1. A large body of literature demonstrates that people tend to keep away from these extremes in their probability judgments, and so are 'conservative' in their probability assessments. These results show that the closer $P(A)$ is to 0, the more $\overline{P_E}(A)$ is greater than $P(A)$, while the closer $P(A)$ is to 1, the more $\overline{P_E}(A)$ is less than $P(A)$. Differences between true and estimated probabilities are low when $P(A)$ is close to 0.5 and increase as $P(A)$ approaches the boundaries of 0 or 1. This pattern was originally seen in research on people's revision of their probablity estimates in the light of further data (Edwards, 1968), and was later found directly in probability estimation tasks. This pattern is sometimes referred to as underconfidence in people's probability estimates (see Erev et al., 1994, for a review).

Conservatism will occur as a straightforward consequence of random variation in our otherwise perfect reasoner. Our reasoner computes $P_E(A)$ by querying episodic memory to count all episodes containing A and dividing by the total number of episodes, giving

$$P_E(A) = \frac{C(A)}{n}$$

Random variation afffects $P_E(A)$ when it causes some flag $f_i(A)$ be read incorrectly. The expected value of $P_E(A)$ is given

$$\overline{P_E}(A) = \frac{T_A(1-d) + (n - T_A)d}{n}$$

(since on average $1-d$ of the T_A flags whose value is 1 will be read as 1, and d of the $n - T_A$ flags whose value is 0 will be read as 1). Since we assume that if the reasoner were not subject to random variation then each estimate $P_E(A)$ would be equal to $P(A)$, we have

$$P(A) = \frac{T_A}{n}$$

and so

$$\overline{P_E}(A) = P(A) + d - 2dP(A) \tag{2}$$

and the average value of $P_E(A)$ deviates from $P(A)$ in a way that systematically depends on $P(A)$. This deviation produces the pattern of conservatism seen in people probability judgments. If $P(A) = 0.5$ this deviation will be 0, if $P(A) < 0.5$ then since d cannot be negative we have $\overline{P_E}(A) > P(A)$ with the difference increasing as $P(A)$ approaches 0, and if $P(A) > 0.5$ then $\overline{P_E}(A) < P(A)$ with the difference increasing as $P(A)$ approaches 1, just as seen in people's probability judgments.

2.2 Subadditivity

Let $A_1 \ldots A_n$ be a set of n mutually exclusive events, and let $A = A_1 \vee \ldots \vee A_n$ be the disjunction (the 'or') of those n events. Then probability theory requires that

$$P(A_1) + \ldots + P(A_n) = P(A)$$

Experimental results show that people reliably violate this requirement, and in a characteristic way. On average the sum of people's probability estimates for events $A_1 \ldots A_n$ is reliably greater than their estimate for the probability of A, with the difference (the degree of 'subadditivity') increases reliably as n increases. An additional, more specific pattern is also seen: for pairs of mutually exclusive events A_1 and A_2 whose probabilities sum to 1 we find that the sums of people's estimates for A_1 and A_2 are normally distributed around 1, and so on average this sum is equal to 1 just as required by probability theory. This pattern is sometimes referred to as 'binary complementarity' (see Tversky and Koehler, 1994, for a detailed review of these results).

Again, these patterns of subadditivity occur as a straightforward consequence of random variation in our otherwise perfect reasoner. From Equation 2 we have

$$\begin{aligned}\overline{P_E}(A_1) + \ldots + \overline{P_E}(A_n) =& P(A_1) + d - 2dP(A_1) + \\ & P(A_2) + d - 2dP(A_2) + \\ & \ldots \\ & P(A_n) + d - 2dP(A_n)\end{aligned}$$

or rearranging

$$\overline{P_E}(A_1)+\ldots+\overline{P_E}(A_n) = (P(A_1) + \ldots + P(A_n)) + nd - 2d\,(P(A_1) + \ldots + P(A_n))$$

Since by definition

$$P(A_1) + \ldots + P(A_n) = P(A)$$

we can rewrite this as

$$\overline{P_E}(A_1) + \ldots + \overline{P_E}(A_n) = P(A) + nd - 2dP(A)$$

Taking the difference between this expression and that for $\overline{P_E}(A)$ (from Equation2) we get

$$\begin{aligned}\overline{P_E}(A_1) + \ldots + \overline{P_E}(A_n) - \overline{P_E}(A) &= P(A) + nd - 2dP(A) - P(A) - d + 2dP(A) \\ &= (n-1)d\end{aligned}$$

and so this difference increases as n increases, producing a pattern of subadditivity just as seen in people's probability judgments. In the case of two mutually excusive events A_1 and A_2 whose probabilities sum to 1, from Equation 2 we get

$$\begin{aligned}\overline{P_E}(A_1) + \overline{P_E}(A_2) &= P(A_1) + d - 2dP(A_1) + P(A_2) + d1 - 2dP(A_2) \\ &= P(A_1) + P(A_2) + d(2 - 2(P(A_1) + P(A_2)) \\ &= P(A_1) + P(A_2) = 1\end{aligned}$$

producing just the pattern of binary complementarity seen in people's probability judgments.

2.3 Conjunction and disjunction errors

The above two biases concern averages of estimated probability values. The next two biases concern differences between probability estimates. Let A_1 and A_2 be any two events ordered so that $P(A_1) \leq P(A_2)$. Then probability theory requires that

$$P(A_1 \wedge A_2) \leq P(A_1)$$

This following from the fact that $A_1 \wedge A_2$ can only occur if A_1 itself occurs. People reliably violate this requirement for some events, giving probability estimates for conjunctions that are greater than the estimates they gave for one or other constituent of that conjunction. This 'conjunction error' does not occur for all or even most conjunctions (people correctly follow the rules of probability theory for most conjunctions). Numerous experimental studies have shown that the occurence of this error depends on the probabilities of A_1 and A_2. In particular, the greater the difference between $P(A_1)$ and $P(A_2)$, the more frequent the conjunction error is, and the greater the conditional probability $P(A_1|A_2)$, the more frequent the conjunction error is. The frequency of the error can be high when these two conditions hold (see Costello, 2009a, for a detailed review).

A similar pattern occurs for people's probability estimates for disjunctions. Since $A_1 \vee A_2$ necessarily occurs if A_2 itself occurs, probability theory requires that

$$P(A_1 \vee A_2) \geq P(A_2)$$

must always hold. People reliably violate this requirement for some events, giving probability estimates for disjunctions that are less than the estimates they gave for one or other constituent. Just as for the conjunction error, the greater the difference between $P(A_1)$ and $P(A_2)$, and the higher the estimated conditional probability $P(A_1|A_2)$, the higher the rate of occurence of the disjunction error. Studies which examine the rate of both errors show a strong correlation between the frequency of the conjunction error for a given pair of events and the frequency of the disjunction error for that same pair (see Costello, 2009b, for a review).

Again, these patterns occur as a straightforward consequence of random variation in our otherwise perfect reasoner. Since the reasoner is subject to random variation we can write

$$\overline{P_E}(A_1) = P(A_1) + d - 2dP(A_1)$$

and

$$\overline{P_E}(A_1 \wedge A_2) = P(A_1 \wedge A_2) + d - 2dP(A_1 \wedge A_2)$$

and we see that the reasoner's estimates for $P(A_1)$ and $P(A_1 \wedge A_2)$ will vary around those means. More specifically the reasoner's estimates for those probabilites at any given moment will be equal to

$$P_E(A_1) = \overline{P_E}(A_1) + e_{A_1}$$

and

$$P_E(A_1 \wedge A_2) = \overline{P_E}(A_1 \wedge A_2) + e_{A_1 \wedge A_2}$$

where e_{A_1} and $e_{A_1 \wedge A_2}$ represent positive or negative random deviation from the mean at that time. The conjunction error will occur when

$$\overline{P_E}(A_1) + e_{A_1} < \overline{P_E}(A_1 \wedge A_2) + e_{A_1 \wedge A_2}$$

or, substituting and rearranging, when

$$(P(A_1) - P(A_1 \wedge A_2))(1 - 2d) < e_{A_1 \wedge A_2} - e_{A_1} \qquad (3)$$

holds. Given that $e_{A_1 \wedge A_2}$ and e_{A_1} vary randomly and can be either positive or negative, this inequality can hold in some cases. The inequality is most likely to hold when $P(A_1) - P(A_1 \wedge A_2)$ is low (because in that situation the left side of the inequality is low). Since

$$P(A_1 \wedge A_2) = P(A_1|A_2)P(A_2)$$

we see that $P(A_1) - P(A_1 \wedge A_2)$ is low when $P(A_1)$ is low and $P(A_1|A_2)$ and $P(A_2)$ are high. We thus expect the conjunction error to be most frequent when $P(A_1)$ is low and $P(A_1|A_2)$ and $P(A_2)$ are both high. This is just the pattern seen when the conjunction error occurs in people's probability estimates.

Reasoning in just the same way for disjunctions, we see that the disjunction error will occur when

$$\overline{P_E}(A_2) + e_{A_2} > \overline{P_E}(A_1 \vee A_2) + e_{A_1 \vee A_2}$$

or, substituting and rearranging as before, when

$$(P(A_1 \vee A_2) - P(A_2))(1 - 2d) < e_{A_2} - e_{A_1 \vee A_2}$$

holds. But from the addition law (Equation 1) we have

$$P(A_1 \vee A_2) - P(A_2) = P(A_1) + P(A_2) - P(A_1 \wedge A_2) - P(A_2) = P(A_1) - P(A_1 \wedge A_2)$$

and substituting we see that the disjunction error will occur when

$$(P(A_1) - P(A_1 \wedge A_2))(1 - 2d) < e_{A_2} - e_{A_1 \vee A_2} \qquad (4)$$

and so, just as with the conjunction error, we expect the disjunction error to be most frequent when $P(A_1)$ is low and $P(A_1|A_2)$ and $P(A_2)$ are both high. More specifically, since the criteria for the conjunction error and the disjunction error are the same (compare the left hand sides of Equations 3 and 4) we expect that on average the rate of occurrence of the disjunction error for a given set of items should match the rate of occurrence of the conjunction error of those items. Again, this is just the pattern seen when the disjunction error occurs in people's probability estimates.

3 Predicting unbiased probability estimates

We saw in the previous section that a simple account where probability estimates are normatively correct but influenced by random noise can explain various patterns of bias in people's probability judgements, and also explain some specific situations in which those biases vanish (when probabilities are close to 0.5, for conservatism; and when two complementary probabilities sum to 1, for subadditivity). We now present a third situation in which this account predicts that bias will disappear.

Consider an experiment where we ask people to estimate, for any pair of events A and B, the probabilities of A, B, $A \wedge B$ and $A \vee B$. For each participant's estimates for each pair of events A and B we can compute a derived sum

$$X_E(A, B) = P_E(A) + P_E(B) - P_E(A \wedge B) - P_E(A \vee B)$$

We can make a specific prediction about the average value of $X_E(A, B)$ for all events A and B. This value will be

$$\overline{X_E}(A, B) = \overline{P_E}(A) + \overline{P_E}(B) - \overline{P_E}(A \wedge B) - \overline{P_E}(A \vee B)$$

From Equation 2 we get

$$\begin{aligned} \overline{X_E}(A, B) =& P(A) + d - 2dP(A) \\ &+ P(B) + d - 2dP(B) \\ &- P(A \wedge B) - d + 2dP(A \wedge B) \\ &- P(A \vee B) - d + 2P(A \vee B) \end{aligned}$$

or rearranging

$$\overline{X_E}(A, B) = (1 - 2d)(P(A) + P(B) - P(A \wedge B) - P(A \vee B))$$

However, from the addition law (Equation 1 we get

$$P(A) + P(B) - P(A \wedge B) - P(A \vee B) = 0$$

for all events A and B, and we see that $\overline{X_E}(A, B) = 0$. Our prediction, therefore, is that on average $X_E(A, B)$ will be equal to 0 for all pairs of events A and B. More specifically, we predict that across a range of judgments for pairs A, B, values of $X_E(A, B)$ will be normally distributed around a mean of 0.

We tested this prediction using data from an experiment on conjunction and disjunction errors carried out by the first author (Experiment 3 in Costello, 2009b). The original aim of this experiment was to examine an attempt by Gigerenzer to 'explain away' the conjunction error as a consquence of people being asked to judge the probability of one-off, unique events. Gigerenzer argued that from a frequentist standpoint the rules of probability theory apply only to repeatable events and not to unique events, and so people's deviation from the rules of probability theory for unique events are not, in fact, fallacious. To assess this argument the experiment examined the occurrence of fallacies in probability judgements for conjunctions and disjunctions of canonical repeatedly-occurring events: weather events such as 'rain','wind' and so on. Contrary to Gigerenzer's argument, participants in these experiments often committed conjunction and disjunction errors; these errors thus cannot be dismissed as an artifact of researchers using unique events in their studies of conjunctive probability.

This experiment gathered probability estimates from 83 participants for A,B, $A \wedge B$ and $A \vee B$ expressions for 12 distinct pairs of events. We briefly present the materials and procedure for this experiment below.

3.1 Materials

Two sets of weather events we used to construct the materials for this experiment: the set 'cloudy','windy','thundery','sunny' and the set 'cold','frosty','sleety'. These sets were selected to contain weather events with a range of different probability values from high (on average participants estimated that 76 days out of 100 are cloudy) to intermediate (participants estimated that 38 days out of 100 are sunny) to low (participants estimated that 12 days out of 100 are thundery).

From these two sets, conjunctive and disjunctive weather events were formed by pairing each member of the first set with every member of the second set and placing 'and'/'or' between the elements of each pair as required. Participants were asked to judge the probability of all single events in the two sets and all conjunctive and disjunctive events constructed from these pairings; there were thus 7 single events, 12 conjunctive events and 12 disjunctive events in total. Each participant gave probability estimates for all these events.

3.2 Procedure

Participants in this experiment were divided into two groups. Participants in the first group were asked questions with the form

- 'What is the probability that the weather will be W on a randomly selected day in Ireland?'

(where W stands for some weather event such as 'sunny', 'windy and sunny', 'windy or sunny', and so on). Participants in the second group was asked questions of the form

- 'Imagine a set of 100 different days, selected at random. On how many of those 100 days do you think the weather in Ireland would be W?'

(where W again stands for some weather event). Participant responses to both frequency and probability rating questions were given on a sliding scale of integers ranging from 0 to 100. Participants answering frequency questions were told that an integer selected on this scale for a given question represented the number of days that participant thought the weather event in question would occur. Participants answering probability questions were told that an integer selected on this scale represented the percentage probability (chance out of 100%)

that the weather event in question would occur. Each participant saw events in a different random order.

3.3 Conjunction and disjunction error rates

A conjunction error was recorded when a participant gave a conjunction a rating five points more probable or more frequent (on the 0 to 100 scale) than one or other constituent. A disjunction error was recorded when a participant gave a disjunction a rating five points less probable or frequent than one or other constituent. While errors occured less often in the frequency group than in the probability group (confirming previous results showing that the frequency format reduces but does not eliminate these errors), there was a significant error rate in both groups (see Costello, 2009b).

Since the distinction between frequency and probability groups is not relevant to our current concerns, we combine the two groups in our analysis. There were 996 distinct conjunction responses in the experiment (83 participants $\times 12$ conjunctions): a conjunction error was recorded in 391 (39%) of those responses. There were also 996 distinct disjunction responses: a disjunction error was recorded in 388 (39%) of these. There were an average of 4.7 conjunction errors and 4.7 disjunction errors per participant; of the 83 participants only 1 did not produce any error responses. There was a significant correlation between the average difference between constituent probability values and the occurence of the conjunction fallacy (r=0.92,p¡0.01) and and between that difference and the occurence of the disjunction fallacy ($r = 0.90, p < 0.01$), just as predicted. There was also a significant correlation between conjunction error rate and disjunction error rate across materials ($r = 0.83, p < 0.01$) showing the predicted relationship between the two patterns of error.

3.4 Values of $X_E(A, B)$

For every pair of weather events A, B used in the experiment, each participant gave probability estimates for the two constituents A and B, for the conjunction $A \wedge B$ and for the disjunction $A \vee B$. Each participant was gave these estimates for 12 such pairs. For each participant we can thus calculate the value $X_E(A, B)$ for 12 pairs A, B, and so across all 83 participants we have 996 distinct values of $X_E(A, B)$. Our prediction is that these values will be normally distributed around 0.

The mean value of $X(A, B)$, averaging across all 996 distinct values obtained in the experiment, was $0.66(SD = 27.1)$. This mean was within 1 unit of the predicted mean on the 100 point scale used in the experiment. Figure 1 shows the raw distribution of these values in the experimental data; Figure 2 gives a simpler picture, showing the distribution of these values in bins of size $v-5 \ldots v+5$ for v from -100 to 100. It is clear from these graphs that $X_E(A, B)$ is normally distributed around this mean (a similar pattern is seen for a range of different bin sizes). The fact that the observed mean differs from the predicted mean by less than seven parts in a thousand is a striking confirmation of the



Figure 1: Scatterplot showing the frequency of different values of $X_E(A,B)$ as observed in the experimental data.

prediction. Using T-tests comparing values of $X_E(A,B)$ with candidate means we obtain a 99% confidence interval of $(-1.6\ldots2.8)$ for the mean. We can thus say with 99% confidence that the population mean for $X_E(A,B)$ is within 2.8 units of the predicted value on the 100 point scale, further confirming the prediction.

4 Discussion

The above result is based on a specific expression $X_E(A,B)$ that cancels out the effect of noise in people's probability judgements. When noise is cancelled in this way we get a mean value for $X_E(A,B)$ that is almost exactly equal to that predicted by probability theory. This close agreement with probability theory occurs alongside reliable and systematic biases towards conjunction and disjunction errors in the same probability judgments. These results provide strong evidence that, when making judgments under uncertainty, people do in fact closely follow the calculus of chance provided by probability theory, and that the observed patterns of bias are due to the systematic influence of noise on those judgements.

Note that we are not suggesting that people are consciously aware of the equations of probability theory when estimating probabilities. That is evidently not the case, given the high rates of conjunction and disjunction errors in people's probability judgments. Indeed we doubt whether any of the participants in our experiment were aware of the invariant required by probability theory's addition law, or would be able to consciously apply that law to their proba-



Figure 2: Scatterplot showing the frequency of different values of $X_E(A, B)$ in bins of size $v - 5 \ldots v + 5$ for v from -100 to 100 in steps of 10.

bility estimations. Instead we propose that people's probability judgments are derived from a 'black box' module of cognition that estimates the probability of an event A by retreiving (some analogue of) a count of instances of A from memory. Such a mechanism is necessarily subject to the requirements of set theory and therefore implicitly embodies the equations of probability theory (which derive from set theory).

We expect this probability estimation module to be unconscious, automatic, rapid, parallel, relatively undemanding of cognitive capacity, and evolutionarily 'old' and so shared between humans and animals. In the terms used by Stanovich and West Stanovich and West (2000), this is a System 1 process, albeit one that is not based on heuristics but instead on the normatively correct rules of probability theory. Support from this modular view comes from that fact that people can make probability judgments rapidly and easily, and typically do not have access to the reasons behind their estimations (using our experiment as an example, if we asked a participant why they judged that rain had 70% probability of occurring, they would not be able to give any answer beyond saying that that was their estimate based on their experience). Further support comes from experimental results showing that animals effectively judge probabilities (for instance, the probability of obtaining food from a given source) and that their judged probabilities are typically close to optimal (for example, see Kheifets and Gallistel, 2012).

We also expect this automatic probability estimation process to be subject to occasional intervention by other cognitive systems, and particularly by conscious, symbolic and slow 'System 2' processes that can check the logical validity of estimates produced by the probability module. We expect this type of intervention to be both relatively rare, and effortful. To quote one participant in an earlier experiment where participants had to choose between betting on a single event or on a conjunction containing that event: ‘I know that the right answer is always to bet on the single one, but sometimes I'm really drawn to the double one, and it's hard to resist’.

4.1 Further predictions

It should be clear that our prediction for the derived expression $X_E(A,B)$ holds because the associated expression $X(A,B)$ sums to 0, and because there are an equal number of positive and negative terms in the expression (these two requirements are necessary to cancel out the d or noise terms in the expression). In this section we give two other expressions where these requirements both hold, and so for which the same prediction follows.

Consider an experiment where we ask people to estimate, for any triplet of events A, B and C, the probabilities of those events and all their conjunctions and disjunctions. As before we can compute a derived sum

$$\begin{aligned} X_E(A,B,C) = P_E(A) + P_E(B) + P_E(C) - P_E(A \wedge B) - P_E(A \wedge C) - P_E(B \wedge C) \\ + P_E(A \wedge B \wedge C) - P_E(A \vee B \vee C) \end{aligned}$$

Applying Equation 2 as before we get

$$\begin{aligned} \overline{X_E}(A,B,C) = &P(A) + d - 2dP(A) \\ &+ P(B) + d - 2dP(B) \\ &+ P(C) + d - 2dP(C) \\ &- P(A \wedge B) - d + 2dP(A \wedge B) \\ &- P(A \wedge C) - d + 2dP(A \wedge C) \\ &- P(B \wedge C) - d + 2dP(B \wedge C) \\ &+ P_E(A \wedge B \wedge C) + d - 2dP_E(A \wedge B \wedge C) \\ &- P_E(A \vee B \vee C) - d + 2dP_E(A \vee B \vee C) \end{aligned}$$

or rearranging

$$\overline{X_E}(A,B) = (1-2d)(P(A)+P(B)+P(C)-P(A\wedge B)-P(A\wedge C)-P(B\wedge C)+P(A\wedge B\wedge C)-P(A\vee B\vee C))$$

However, as before probability theory's addition law gives

$$P(A)+P(B)+P(C)-P(A\wedge B)-P(A\wedge C)-P(B\wedge C)+P(A\wedge B\wedge C)-P(A\vee B\vee C) = 0$$

for all events A, B and C, and again we see that

$$\overline{X_E}(A,B,C) = 0$$

will hold for all triples of events A, B, C, and so we predict that across a range of participants and event triples the value of $X_E(A,B,C)$ will be normally distributed around a mean of 0.

Next consider an experiment where we ask people to estimate, for any pair of events A and B, the probabilities of A, B, $A\neg B$ ('A and not B') and $B\neg A$ ('B and not A') . In this case our derived sum is

$$Y_E(A,B) = P_E(A) + P_E(B) - P_E(A\neg B) - P_E(B\neg A)$$

whose expected value will be

$$\overline{Y_E}(A,B) = \overline{P_E}(A) + \overline{P_E}(A\neg B) - \overline{P_E}(B) - \overline{P_E}(B\neg A)$$

From Equation 2 we get

$$\begin{aligned}\overline{Y_E}(A,B) =& P(A) + d - 2dP(A) \\ &+ P(A\neg B) + d - 2dP(A\neg B) \\ &- P(B) - d + 2dP(B) \\ &- P(B\neg A) - d + 2P(B\neg A)\end{aligned}$$

or rearranging

$$\overline{Y_E}(A,B) = (1-2d)(P(A) + P(A\neg B) - P(B) - P(B\neg A))$$

However, from probability theory we have

$$P(A) + P(A\neg B) = P(B) + P(B\neg A)$$

(because each side of the expression is equal to $P(A \vee B)$) and so we have

$$P(A) + P(A\neg B) - P(B) - P(B\neg A) = 0$$

for all events A and B, and again we predict that across a range of participants and event triples the value of $Y_E(A,B)$ will be normally distributed around a mean of 0.

We could, of course, run experiments testing these two predictions ourselves. Given the strong confirmation of our original prediction, however, we are convinced that these predictions will be confirmed: for us, running those experiments would in some ways be a waste of time. We feel it would be more useful for others to test these predictions. We hope that other researchers take up this challenge and gather further evidence that people's mechanisms for judging probability are not systematically flawed, but instead are based the rules of probability theory (but subject to the influence of noise).

References

E. Brunswik. In defense of probabilistic functionalism: A reply. *Psychological Review*, 62:236–242, 1955.

F. Costello. How probability theory explains the conjunction fallacy. *Journal of Behavioral Decision Making*, 22(3):213–234, 2009a.

Fintan J. Costello. Fallacies in probability judgments for conjunctions and disjunctions of everyday events. *Journal of Behavioral Decision Making*, 22 (3):235–251, 2009b.

W. Edwards. Conservatism in human information processing. In B. Kleinmuntz, editor, *Formal Representation of Human Judgment.*, pages 17–52. New York: Wiley, 1968.

Ido Erev, Thomas S. Wallsten, and David V. Budescu. Simultaneous over- and underconfidence: The role of error in judgment processes. *Psychological Review*, 101(3):519–527, 1994.

M. Hilbert. Toward a synthesis of cognitive biases: how noisy information processing can bias human decision making. *Psychological Bulletin*, 138(2): 211–237, 2012.

A. Kheifets and C. R. Gallistel. Mice take calculated risks. *Proceedings of the National Academy of Sciences*, 2012. doi: 10.1073/pnas.1205131109. URL `http://dx.doi.org/10.1073/pnas.1205131109`.

Eldar Shafir and Robyn A. Leboeuf. Rationality. *Annual Review of Psychology*, 53(1):491–517, 2002.

Keith E. Stanovich and Richard F. West. Individual differences in reasoning: Implications for the rationality debate? *Behavioral and Brain Sciences*, 23 (5):645–665, 2000.

A. Tversky and D.J. Koehler. Support theory: A nonextensional representation of subjective probability. *Psychological Review*, 101(4):547–566, 1994.

Amos Tversky and Daniel Kahneman. Availability: A heuristic for judging frequency and probability. *Cognitive Psychology*, 5:207–232, 1973.